Scanning For Interference Fields

Key Areas to Scan to Identify Interference Fields

An interference field is an area of the body (small or large) that tests abnormally as a result of individual or multiple factors, such as scars, physical or emotional trauma, infection, nutritional deficiencies, heavy metal toxicity, etc. An interference field can impede the normal, healthy electromagnetic flow of the body's acupuncture meridian system, thus creating symptomology that is difficult to trace back to its original root cause by conventional methods. However, identifying an interference field can now be easily done using the Bi-Digital O-Ring Test (as developed by Dr. Yoshiaki Omura, MD, Sc.D.)

Note: the size of an interference field does not dictate its importance. For example: larger scars are generally more troublesome, but a very small scar can also be highly reactive and create troublesome symptoms.

Keep in mind: in general, the patient often does NOT complain about the site of the interference field. It is usually the reflex area of the interference field that creates the distressing symptoms.

Examples of Common Interference Fields

Scars

Areas where the epidermis of the skin has been cut

- Any type of surgical scars
  - Scars from invasive procedures (such as those for the gallbladder, the heart or for cancer)
  - Mole or skin cancer removal scars
  - Episiotomy scars, C-section scars, breast implants, tummy tucks, cosmetic surgery
  - Shunt scars
- Internal scars
  - Scars that may not be visible: internal hysterectomy scars, internal sinus surgery, etc.
  - Exploratory surgery
  - Hernia repair
- Traumatic scars: from falls, cuts, etc.
- Puncture areas: from hypodermic injections, intravenous Drug Therapy, spinal blocks, etc.
- Acne scars; chicken pox scars, animal bites, bites from mosquitos or other insects, etc.



86-Year-Old Woman
C-section scar (from 55 years ago) blocking 8 different meridians, reflexing downwards to cause stiff, swollen legs, intractable to therapy



30-Year-Old Businessman
Vaccination scar (from 20 years ago) blocking outflow of head and neck meridians, creating stiff neck, not resolved inspite of many therapies



42-Year-Old Housewife
Episiotomy scar (from 15 years ago) reflexing to left leg creating sciatica pain which would not resolve with many types of therapies

© Copyright 2004 Premier Research Labs

- Vaccination scars

Traumatized Areas

Previous areas of trauma where there are no visible scars

- Trauma from car accidents: whiplash, sprains/strains, smashed areas, etc.
- Trauma from other sources: falls, athletic sprains, strains, scrapes, bumping head, etc.
- Impact sites: (i.e. an area previously hit with a fast ball, etc.)
- Burned areas (scars or no scars), skin grafts
- Numb areas



18-Year-Old Student
Weak neck area from previous car accident (5 years ago) reflexing to liver area to create allergic reactions



55-Year-Old Sales Director
Weak area from previous fall on hip area (15 years ago) reflexing to abdominal area creating excess abdominal weight



45-Year-Old Journalist
Weak area due to ankle sprain (10 years ago) reflexing to heart area, creating heart palpitations

Abnormal Body Areas

Areas that look or feel abnormal

- Bruises or discolored skin areas that do not heal in a reasonable amount of time
- Skin rashes, nail fungus or sores that do not heal in a reasonable amount of time
- Areas that itch or have abnormal sensations: i.e. burning, cold, tingling, etc.



58-Year-Old Female Caretaker
Bruise on left leg area after a fall (1 year ago) still there, reflexing to abdominal area creating weight gain



67-Year-Old Male CEO
Toe nail fungus on both feet (over 30 years) reflexing to stomach area to create excess stomach weight



27-Year-Old Male Mechanic
Decaying 3rd molar reflexing to chest to create a jittery feeling that then creates insomnia at night

© Copyright 2004 Premier Research Labs

Permanent Tattoos



50-Year-Old Female Entertainer
Bilateral buttocks tattoos reflexing to various organs to create chronic fatigue



22-Year-Old Male Marketeer
Tattoo on upper right arm reflexing to right side of neck, causing neck stiffness



40-Year-Old Female Executive
Tattoo on right inner ankle reflexing to kidney to create tiredness

Ear and Body Piercings



35-Year-Old Female Receptionist
Pierced earlobes reflexing to stomach, causing weight gain and indigestion



21-Year-Old Male Student
Metal tongue piercing reflexing to head area, causing intermittent dizziness and fatigue



25-Year-Old Medical Receptionist
Metal naval piercing reflexing to throat, making it difficult to swallow.

Metal Inside the Body

- Surgical metal pins, plates, mesh; bullet or shrapnel fragments
- Dental metal: silver fillings, god crowns, stainless steel posts, composite fillings (typically high in aluminum), dental implants (typically titanium)



30-Year-Old Horse Trainer
Metal plate in head after car accident reflexing to head and kidneys, creating headaches and fatigue



45-Year-Old Housewife
Staples from previous bladder surgery (8 years ago) reflexing to lower back, creating constant back pain



60-Year-Old Stock Trader
Gold crowns blocking meridian outflow of mouth reflexing to multiple meridians, causing weight gain, intestinal gas and sleep disturbances

Injections

- Dental Injections. Injection sites in the gum areas where dental anesthetics have been used (especially retromolar areas).
- Neural Therapy. Injection sites where anesthetics (Procaine, Carbocaine) have been injected into the body, including dental areas.
- IV Drips. Injection sites where intravenous needles were placed in the arm.
- Other Injections. Injection sites where drugs, vitamins (such as B-12 shots) or other subsances have been injected into the body.

Dental Injections



45-Year-Old Writer
Previous dental injections in the retromolar areas blocking the energetic outflow to the jaw, creating a chronic stiff neck

Neural Therapy



50-Year-Old Medical Doctor
Previous injection sites with Procaine in navel area reflexing to stomach and head area, causing poor digestion and headache

IV Therapy



40-Year-Old Tennis Player
Repeated IV needle injections into right arm reflexing to back of neck, causing fatigue

Dental and Gum Areas

- Hidden dental decay under silver fillings, composites or crowns
- Toxins exuding from conventional root canals



75-Year-Old Retired CEO

Lower lateral front tooth with decay reflexing to heart, causing heart palpitations and shortness of breath.



62-Year-Old Restaurant Owner

Conventional root canal on front tooth reflexing to prostate, causing prostate cancer



42-Year-Old Housewife

Decayed upper left molar reflexing to left ear, causing chronic ear infection and pain

The SPLAT Effect

One injury site can sedate multiple body sites.

When an injury occurs, if the body is nutritionally deficient or immune-compromised at the time of the injury injury site can become an interference field which routinely reflexes to one or more areas of the body — w can be small, medium or large reflex areas — creating entrenched symptoms. We call this the Splat Effect cause one injury site can reflex like a splat to many areas.



The Bio-Electric Body

By using QRA™ to test the various field-sedating reflex areas (also called "referred areas") arising from an injured body site clearly demonstrates the bio-electric nature of the body.

DOUBLE JEOPARDY

Multiple Interference Fields Reflexing to One Area

Frequently, one or more interference fields may reflex to the same body area. In these cases, the symptoms will not abate until all the interference fields have been cleared. In cases like these, some call it double jeopardy (i.e. 2 interference fields are reflexing to the same area) or triple jeopardy (i.e. 3 interference fields are reflexing to the same area) or quadruple jeopardy, etc. In these cases, clearing one interference field will help improve the symptom area but the symptom may not be completely and permanently cleared until all interference fields have been eliminated.

Symptom: Chronic Right Knee Pain

QRA™ Analysis: Four interference fields all reflexing to the same knee area



Rev. 02-18-08

Frequently Overlooked Interference Fields

Of course, the majority of a practitioner's time must be spent tracking down interference fields that are directly related to their patient's main presenting symptoms. However, the following areas are often interference fields that may not be directly related to the patient's chief complaints, but nonetheless, may be contributing to the body's overall burden of bioenergetic dysfunction.

Therefore, after a significant amount of progress has been made in clearing the patient's chief complaints, the following areas should be checked and treated if testing weak:

- **Fingernails and Toenails** (esp. women) - These areas often test toxic due to the use of toxic commercial fingernail and toe polish (including so-called "natural" ones). Even after using Medi-Body Packs to clear the top and bottom of the feet (so they now test strong) and the palms and back of the hands (so they now test strong) – often, the nails themselves will still test weak due to toxic chemical absorption from commercial polishes – *no matter how long ago in the past they were used.* Recommend discontinuing all commercial finger and toenail polishes.

- **Around eyes and eyelids** (esp. women) – These areas frequently test toxic due to the routine use of toxic eye cosmetics. Be sure to test these areas, then treat and clear the areas that test weak.

- **Lips (esp. women)** – The lips frequently test toxic due to the frequent use of toxic lipstick and lip balms (yes, even so-called "organic" ones). Be sure to test the lips, upper and lower, then treat and clear the areas that test weak.

- **Eyebrows (esp. women)** — These areas frequently test toxic due to the frequent use of toxic eyebrow cosmetics (even so-called "organic" ones). Test each eyebrow, then treat and clear the areas that test weak.

- **Navel** - The navel is the usually the first scar in life for most people. If, during the birthing process, the umbilicus has been cut too soon (very common), then the navel area will test weak (*often very weak*), thus weakening the bioenergetic flow of the main central meridian on the front of the body (the CV channel) – even from birth. Be sure to test the navel area, then treat and clear it if it tests weak.

- **Pierced ears** – This area frequently tests weak due to puncturing the ear at a time when the person had insufficient nutrients to adequately heal the ear scar or due to being ill during that time. The vast majority of body piercings we have tested were weak. Be sure to test all body piercings, then treat and clear any that test weak.

- **Tattooes** -- These areas typically test weak and sedate the meridians where they have been injected (often creating distant, distressing symptoms not usually traced back to the tattoo interference field). By using Medi-Body Packs, over time these areas will gradually begin to test strong again. Be sure to test every tattoo (no matter now small), then treat and clear each one that tests weak. Recommend the use of "temporary" tattooes (if desired) that can be washed off or peeled off.

- **Scalp (esp. women)** – Many areas of the scalp frequently test weak due to the use of toxic bleaches, hair dyes and/or hair permanent solutions. It is important to treat and clear every area of the scalp that tests weak.

- **Perfume or Afteshave** — Most commercial perfumes typically contain a myriad of untested, ingredients which are almost all toxic. If a person has used commercial perfumes, colognes or aftershave – *no matter how long ago in the past* – then test each area (such as around the neck). If any of these areas test weak, then treat each one until it tests clear. Recommend natural, quantum-state essential oils as an excellent daily, health-promoting perfume.

- **Armpits** -- If a person has used aluminum-based deodorants or other toxic commercial deodorants – *no matter how long ago in the past*— then test both armpit areas. If either or both test toxic, then treat them until they test clear.

- **Vaginal Area** – If a woman has frequently used spermacide lotions (such as nonoxynol-9), often the vaginal area will test weak. If so, treat this area until it tests clear. Longterm use of toxic spermacides can contribute to vaginal prolapse later on – or even cancer.

- **Sunscreen** -- If a person has frequently used commercial sunscreen lotions – *no matter how long ago in the past* – test and treat the target skin areas until they test clear. Many commercial sunscreens contain highly questionable ingredients, often including known carcinogens (including so-called "natural" ones) – unfortunately, these toxic elements can be easily absorbed by the skin.

- **Hair Plugs** -- Many balding men, in an effort to look younger, may have had artificial "plugs" of hair surgically implanted in their scalp (often near the front hairline). These areas almost always test weak and should be treated until they test strong.

- **Circumcision Scar** – The scar left from removing the foreskin on the penis frequently tests off (*often very weak*). If it tests weak, this tends to sedate and weaken the main central meridian on the front of the body (the CV meridian). Test and treat the area if it tests weak. Some researchers believe that circumcision may be an unrecognized, contributing factor in the high rates of impotence in older American men. (Could this be due to the blocking effects of the penile scar?)

- **Laser Eye Surgery**. Laser eye surgery can create interference fields in the eyes that often reflex downward from the eyes (to the stomach and abdomen areas). We have noted that the reflexes often create digestive problems later on. Buyer, beware!

Testing Sensitive Areas. If a male practitioner or female practitioner is testing a sensitive body area (such as the vaginal area, a circumcision scar or a tattoo in an unusual place), it is best to discuss the matter ahead of time with the patient to obtain their consent and to help them understand the importance of testing these areas. In addition, even though the patient is fully clothed, it is best to have an assistant in the room during the testing.